

2602 S 24th St, Phoenix, AZ 85034

Tel 602.759.1245 Fax 602.759.8331

www.CyrexLabs.com



PATIENT Name: DOB: Gender:

ACCESSION #: **REQUISITION #: SAMPLE TYPE:** DOCTOR/PATIENT ID: DATE COLLECTED: DATE RECEIVED: DATE OF REPORT:

TEST	RESULTS			
Array 5 - Multiple Autoimmune Reactivity Screen **	IN RANGE (Normal)	EQUIVOCAL*	OUT OF RANGE	REFERENCE RANGE (ELISA Index)
Parietal Cell + ATPase	0.29			0.0 - 2.21
Intrinsic Factor	1.43			0.0 - 2.51
ASCA + ANCA	0.46			0.4 - 1.61
Tropomyosin	1.13			0.0 - 3.01
21-Hydroxylase (Adrenal Cortex)	0.60			0.6 - 1.91
Myocardial Peptide			3.87	0.0 - 1.91
Alpha-Myosin	0.42			0.6 - 2.01
Phospholipid	0.58			0.4 - 1.71
Platelet Glycoprotein	0.32			0.6 - 1.81
Ovary/Testis ***	0.58			0.6 - 1.81
Fibulin	0.30			0.4 - 1.31
Collagen Complex	0.77			0.0 - 1.41
Arthritic Peptide	0.70			0.0 - 1.71
Osteocyte	1.34			0.7 - 2.01
Cytochrome P450 (Hepatocyte)	0.49			0.8 - 2.31
Myelin Basic Protein	0.36			0.6 - 1.71
Asialoganglioside	0.85			0.3 - 2.01
Alpha-Tubulin + Beta-Tubulin	0.64			0.0 - 2.71
Cerebellar	0.45			0.4 - 1.51
Synapsin	1.68			0.0 - 2.11

TEST	RESULTS				
Array 5 - Multiple Autoimmune Reactivity Screen **	NEGATIVE	WEAK POSITIVE	MODERATE TO STRONG POSITIVE	REFERENCE (WHO Units)	
Thyroglobulin IgG	11.84		3 I KONG POSITIVE	0.0 - 200.00	

TEST Array 5 - Multiple Autoimmune Reactivity Screen **	RESULTS NEGATIVE	POSITIVE	REFERENCE	UNITS
Thyroid Peroxidase IgG	59.99		0.0 - 100.00	WHO Units
IA-2 Autoantibody	<0.75		0.0 - 7.49	U/mL
Glutamic Acid Decarboxylase Autoantibody	<5.00		0.0 - 5.00	IU/mL



2602 S 24th St, Phoenix, AZ 85034

Tel 602.759.1245 Fax 602.759.8331

www.CyrexLabs.com



TITIONER Nai

Name:
DOB:
Gender:

ACCESSION #:
REQUISITION #:
SAMPLE TYPE:
DOCTOR/PATIENT ID:
DATE COLLECTED:
DATE RECEIVED:
DATE OF REPORT:

\* Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results >1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.

\*\* All analytes, except for Thyroid Peroxidase, Thyroglobulin, GAD, and IA-2 are tested for IgG and IgA combined. Thyroid Peroxidase (TPO) and Thyroglobulin (Tg) are tested for IgG only. GAD and IA-2 are tested for multiple autoantibodies. TPO results were obtained with the Inova Diagnostics QUANTA Lite TPO ELISA Kit. Thyroglobulin results were obtained with the Inova Diagnostics QUANTA Lite Thyroid T ELISA Kit. GAD65 results were obtained with the Kronus Glutamic Acid Decarboxylase (GAD) Autoantibody ELISA Kit. IA-2 results were obtained with the Kronus IA-2 Autoantibody (IA-2Ab) ELISA Kit. Values obtained with different manufacturers' assay methods may not be used interchangeably. The magnitude of the reported IgG levels cannot be correlated to endpoint titer. Shipping and storage criteria for specimens may vary from Inova Diagnositics' and Kronus IA-2 protocol, however, this does not appear to significantly affect patient results.

\*\*\* Ovary and Testis are tested together to avoid any confusion arising out of potential cross-reactivity.

Sadi Koksoy, DVM, PHD, HCLD(ABB), Laboratory Director

Cyrex Laboratories is certified under the Clinical Laboratory Improvement Amendments of 1988 ("CLIA") as qualified to perform high-complexity clinical testing. Test result data on its own does not constitute a diagnosis of any disease. Only a physician or qualified healthcare professional should interpret the significance of a clinical lab test or make a diagnosis. This test was developed and its performance characteristics determined by Cyrex Laboratories, LLC. This test is a laboratory developed test and therefore not subject to clearance or approval by the US Food and Drug Administration. The names and titles of tests and arrays are for reference purposes only.